



# Modelling temperature-dependent bionomics of *Bemisia tabaci* (Q-biotype)

Olivier Bonato, Amandine Lurette, Claire Vidal, J. Fargues

## ► To cite this version:

Olivier Bonato, Amandine Lurette, Claire Vidal, J. Fargues. Modelling temperature-dependent bionomics of *Bemisia tabaci* (Q-biotype). *Physiological Entomology*, 2007, 32 (1), pp.50-55. 10.1111/j.1365-3032.2006.00540.x . ird-01224737

**HAL Id: ird-01224737**

**<https://hal.ird.fr/ird-01224737>**

Submitted on 5 Nov 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1    **Modelling temperature-dependent bionomics of *Bemisia tabaci* (Q-**  
2    **biotype)**

3  
4    OLIVIER BONATO<sup>1</sup>, AMANDINE LURETTE, CLAIRE VIDAL and JACQUES  
5    FARGUES

6    IRD-INRA Centre for Biology and Management of Populations (CBGP), Montferrier /  
7    Lez cedex, France

8  
9    **Abstract**

10    The influence of temperature (17, 21, 25, 30 and 35°C) on life history traits of a Q-  
11    biotype *Bemisia tabaci* population on tomato is studied. Temperature dependent  
12    relationships are characterized for immature developmental rate, immature survival,  
13    fecundity, longevity and intrinsic rate of increase. Development time vary from 20  
14    days at 30°C to 56 days at 17°C and the lowest thermal threshold is estimated at  
15    10.2°C. The optimal temperature for immature development is 32.5°C. Total  
16    fecundity (eggs per female) ranges from 105.3 (at 21°C) to 41 (at 35°C). The  
17    longevity decreases with temperature increase. The intrinsic rate of increase ranges  
18    from 0.0450 (at 17°C) to 0.123 (at 30°C). The functional relationships between  
19    temperature and life-history parameters are used to evaluate the effect of temperature  
20    on the population dynamics. Such mathematical relationships could provide a basis  
21    for future development of population models.

22  
23    **Key words**

---

<sup>1</sup> Correspondence : Dr Olivier Bonato, IRD-CBGP, CS 30 016, 34988 Montferrier/Lez, France. Tel. :  
+33 467 99 623306 ; fax : +33 467 99 623345 ; e-mail : bonato@mpl.ird.fr

24 Temperature, bionomics, *Bemisia tabaci*, Q biotype, modelling

25

## 26 **Introduction**

27

28 The sweetpotato whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) is  
29 one of the most serious agricultural pest on tomato *Lycopersicum esculentum* (Mill)  
30 and other horticultural crops in tropical and subtropical temperature regions world-  
31 wide. Damage may be caused directly by feeding on phloem or deposition of  
32 honeydew, or indirectly by transmitting different types of plant viruses, such as the  
33 tomato yellow leaf curl virus (Oliveira *et al.*, 2001), to a wide range of vegetable  
34 crops. The potential of *B. tabaci* to develop resistance in response to intensive use of  
35 pesticides has led to studies on integrated pest management strategies in which  
36 biological control plays a central role, and significant advances have been made in  
37 developing and implementing management systems (Gerling & Mayer, 1996;  
38 Naranjo, 2001). As a general rule, any pest management programme should be based  
39 on adequate knowledge of the main factors responsible for changes in population  
40 dynamics. Life history parameters estimated under different biotic or abiotic  
41 conditions provide the basic tools for among others, understanding changes in the  
42 status of pest species (Poole, 1974; Dempster, 1975; Krebs, 1978; Southwood, 1978).  
43 Because *B. tabaci* is a poikilothermic organism i.e. temperature influences the life  
44 table components, it is important to take this factor into consideration in explaining  
45 population ecology. A review by Drost *et al.* (1998) reported that biological  
46 parameters of *B. tabaci* have been characterized for different temperatures, host  
47 plants and biotypes. Among this vast amount of literature, the most complete work is  
48 probably that of Wang & Tsai (1996) concerning B-biotype reared on aubergine.

49 The Q-biotype of *B. tabaci* was first characterized in samples collected in the south  
50 of Spain and Portugal (Guiro *et al.*, 1997). Successive surveys showed that this  
51 biotype is also present in Tunisia (Chermitti *et al.*, 1997), Morocco (Monci *et al.*,  
52 2000), Egypt (De Barro *et al.*, 2000), Israel (Horowitz *et al.* 2003) and southern Italy  
53 (Demichelis *et al.*, 2000; Simón *et al.*, 2003). Because of its high degree of  
54 polyphagy and its ability to transmit a relatively wide range of plant viruses, the Q-  
55 biotype is considered as a particularly dangerous biotype (Muñiz, 2000; Navas-  
56 Castillo *et al.*, 2000). In spite of the wide distribution of the Q-biotype in the  
57 Mediterranean basin, no complete published work is available currently on its life-  
58 history parameters in relation to temperature when reared on tomato. The aim of the  
59 present study was to characterize and analyze functional relationships between  
60 temperature and life-history parameters and to evaluate the effect of temperature on  
61 the dynamics of Q-biotype populations.

62

## 63 **Materials and methods**

64

### 65 *Whitefly source and host plant production*

66

67 In 2002, founders of *Bemisia tabaci* of the Q biotype were collected from a  
68 greenhouse of tomato crop located in Alenya in the South of France (42°38'N;  
69 2°58'E). The stock colony was reared and maintained on tomato plants of the cv  
70 Hilario® (Royal Sluis, Enkhuizen, The Netherlands) in plastic cages placed in  
71 climatic chambers at  $25 \pm 1^\circ\text{C}$ , and  $60 \pm 5\%$  RH. Experiments were carried out using  
72 progeny after more than 3 generations on Hilario.

73

#### 74 *Development of immatures*

75

76 Young adults of *B. tabaci* (150 pairs: male and female) were placed in cubic  
 77 screened cages (50 x 50 x 50cm) each containing a young potted tomato plant. The  
 78 cages containing adults were maintained in a growth chamber at  $25 \pm 1^\circ\text{C}$ ,  $60 \pm 5\%$   
 79 RH in a LD 14:10 h photocycle. Whitefly adults were given 3 h to lay eggs. The  
 80 adults were then removed, leaves were observed under a stereo-microscope at 36X  
 81 magnification. Two eggs on the abaxial surface of 5 leaves were kept and excess  
 82 eggs were killed using an insect pin. Plants were then placed in growth chambers set  
 83 at 5 constant temperatures: 17, 21, 25, 30 and  $35^\circ\text{C}$  with six replicates. Once the eggs  
 84 hatched and the crawlers fixed on the leaf, young nymphs were identified  
 85 individually. Each nymph was observed daily until adult emergence, and the  
 86 transition from one stage to another was noted. Differences between developmental  
 87 times were tested by one-way ANOVA and means were separated by Newman–  
 88 Keuls test ( $\alpha=0.05$ ). Statistical analyses were performed using XLStat 7.1  
 89 (Addinsoft). The influence of temperature ( $T$ ) on developmental rate was described  
 90 by the model proposed by Logan *et al.* (1976):

$$91 \quad DR = p1 \times \left[ \exp(p2 \times (T - Ti)) - \exp\left(p2 \times (Tm - Ti) - \left(\frac{1}{p3}\right) \times (Tm - T)\right) \right]$$

92 where  $DR$  is the development rate which is the reciprocal of development time,  $Ti$  is  
 93 the lower temperature tested and  $Tm$  is the upper threshold derived from the  
 94 observations. The parameters,  $p1$ ,  $p2$  and  $p3$  were estimated by regression. The  
 95 lowest thermal threshold for development ( $LTT$ ) was calculated by the ratio:  $LTT =$   
 96  $a/b$ ,  $a$  and  $b$  were determined by linear regression of the equation  $DR = a + bT$ , for  
 97 temperature ( $T$ ) interval over which the relation was linear.

98 Differences between survival rates were tested using  $\chi^2$  test ( $\alpha=0.05$ ). The  
 99 relationship between temperature ( $T$ ) and immature survival rate was described by  
 100 the Curry & Feldman (1987) model:

101

$$SI = \frac{\left( (T + 273) \times \exp \left( p1 - \left( \frac{p2}{(T + 273)} \right) \right) \right)}{\left( 1 + \exp \left( p3 + \left( \frac{p4}{(T + 273)} \right) \right) \right) + \exp \left( p5 - \left( \frac{p6}{(T + 273)} \right) \right)}$$

105

106 Where  $SI$  is survival of immatures,  $T$  is temperature in °C, and  $p1$  to  $p6$  are  
 107 regression coefficients.

108

#### 109 *Reproductive capacity and female longevity*

110

111 One newly emerged (<24 h) female and two males were placed in a clip-cage on the  
 112 abaxial surface of new leaflets. For each temperature tested, 30 clip-cages were kept  
 113 in growth chambers ( $60 \pm 1\%$  RH, LD 14:10h photocycle). The clip-cages with  
 114 insects were moved to new leaves daily and the number of eggs laid per female was  
 115 counted until death of the female. Differences in fecundity and female longevity  
 116 were compared with ANOVA followed by a Newman–Keuls tests ( $\alpha=0.05$ ).  
 117 Exponential functions were used to describe the influence of temperature on total  
 118 fecundity and longevity.

119 Fecundity:

120

$$EN = p1 \times \left[ (T^{p2}) \times \exp (p3 \times T) \right]$$

121

122 Where  $EN$  is total number of eggs laid per female;  $T$  is temperature in °C; and  $p1$ ,  
123  $p2$ ,  $p3$  are regression coefficients.

124

125 Longevity:

$$126 \quad L = \exp [p1 + p2 \times T]$$

127

128 Where  $L$  is longevity per female in days;  $T$  is temperature in °C; and  $p1$ ,  $p2$ : are  
129 regression coefficients.

130

131 *Demographic parameters*

132

133 The net reproductive rate ( $R_0$ ), the mean generation time ( $G$ ), the intrinsic rate of  
134 natural increase ( $r_m$ ) and the finite rate of increase ( $\lambda$ ) were determined using the  
135 program developed by Hulting *et al.* (1990) in which the parameters are calculated  
136 using the method recommended by Birch (1948). The program, based on Jackknife's  
137 procedures, gives a variance and hence a standard error to each parameter calculated  
138 enabling statistical comparison of values (Meyer, 1986). The relationship between  $r_m$   
139 and temperature was also described using the Logan *et al.* (1976) model (see above).

140

## 141 **Results**

142

143 *Development of immatures*

144

145 *B. tabaci* required 56 days at 17°C to complete its development from egg to adult,  
146 but only 20 days at 35°C (Table 1). Between 17 and 30°C, the developmental time  
147 was negatively correlated with temperature, but no significant difference was found  
148 between 30 and 35°C ( $P>0.05$ ). Values of Logan equation parameters were  $p1 =$   
149  $0.0115$ ,  $p2 = 0.0921$ ,  $p3 = 0.3133$  ( $R^2= 0.92$ ,  $P<0.01$ ). Between 17 and 30°C, the  
150 relation was linear and the *LTT* was estimated at 10.2°C (Fig. 1). The optimal  
151 temperature for development, calculated from the derivative Logan equation, was  
152 32.5°C.

153 Survival rates at 21, 25 and 30°C (Table 2) were not significantly different  
154 ( $\chi_w^2=0.695$ ,  $P=0.707$ ) even if the highest percentage was measured at 25°C.  
155 Temperature of 17°C had the greatest effect on immature development especially in  
156 the 4<sup>th</sup> stadium. The curve obtained by fitting the Curry & Feldman model to the data  
157 ( $p1= -6.55$ ,  $p2= -200.18$ ,  $p3= -2986.08$ ,  $p4= 865870$ ,  $p5= 1353.19$ ,  $p6= 6417118$ ,  
158  $R^2=0.99$ ,  $P <0.01$ ) described the influence of temperature very well and indicated that  
159 17 and 35°C were close to the lower and upper thermal limits, respectively (Fig. 2).

160

#### 161 *Longevity and reproductive capacity of females*

162

163 Temperature had a significant effect on the fecundity and longevity of females.  
164 Longevity was negatively correlated with temperature. The longest life was recorded  
165 at 17°C and the shortest at 35°C (Table 3). The pre-oviposition period was very short  
166 (less than 24h) for almost all females tested and oviposition period was close to  
167 longevity for the 5 temperatures tested. The relation between temperature and female  
168 longevity (Fig. 3) was very well described by the exponential function ( $p1= 5.02$ ,  
169  $p2=-0.079$ ,  $R^2=0.99$ ,  $P <0.01$ ).



170 Except for 17°C, fecundity followed the same trend as longevity, i. e., was negatively  
171 correlated with temperature. Figure 4 shows the influence of temperature on  
172 fecundity ( $p1= 9.9\text{e-}11$ ,  $p2=12.8$ ,  $p3=0.54$ ,  $R^2=0.89$ ,  $P<0.01$ ). Our results showed  
173 that optimal fecundity was obtained from temperatures ranging between 21 and  
174 25°C. The sex-ratio calculated for each treatment was not significantly different than  
175 50% ( $P>0.05$ ).

176

#### 177 *Demographic parameters*

178

179 Parameters were calculated with the sex-ratio of *B. tabaci* set to 0.5. At the five  
180 temperatures tested, the highest rate of increase ( $r_m$ ) was obtained at 30°C and the  
181 lowest at 17°C (Table 4). At 25, 30 and 35°C, the rates of increase were not  
182 significantly different ( $P>0.05$ ). The net reproductive rate was lower at 17°C and  
183 35°C indicating the proximity of lower and upper thermal thresholds, respectively.  
184 The curve (Fig.5) obtained by fitting the Logan model ( $p1 = 0.019$ ,  $p2 = 0.099$ ,  $p3 =$   
185  $0.331$ ,  $R^2= 0.99$ ,  $P<0.01$ ) describes the influence of temperature on the rate of  
186 natural increase very well. The optimal temperature for population development,  
187 calculated from the derivative Logan equation was 31.3°C.

#### 188 **Discussion**

189 Based on biological and ecological information published in the two last decades, the  
190 immature developmental time of *B. tabaci* (from egg to adult) depends on the host  
191 plant (Coudriet *et al.* 1985; Van Lenteren & Noldus, 1990; Bethke *et al.*, 1991;  
192 Zalom *et al.*, 1995; Tsai & Wang, 1996; Muñiz & Nombela, 1997; Nava-Camberos  
193 *et al.*, 2001) as well as on the whitefly populations or biotypes (Drost *et al.*, 1998;  
194 Muñiz, 2000; Muñiz & Nombela, 2001). The developmental time of *B. tabaci*

195 recorded at 25°C ranges from 17.3 to 22.8 days when reared on either aubergine,  
196 tomato, sweet potato, cucumber, bean or pepper. The present results show that on  
197 tomato, the population of Q-biotype *B. tabaci* newly introduced in southern France  
198 require 25.6 days to complete development from egg to adult at 25°C. In contrast,  
199 Tsai & Wang (1996) report a shorter developmental time (17.96 days) for a Floridian  
200 *B. argentifolii* population on tomato (cultivar Suny Hybrid) at 25°C. Based on the  
201 model of Logan *et al.* (1976), the optimal temperature for the development of  
202 immatures (32.5°C) is higher than that calculated for Florida, Mississippi, and Arizona  
203 populations of *B. argentifolii* (29.9, 28.2, and 30.0°C, respectively) on aubergine  
204 (Wang & Tsai, 1996). The optimum for *B. tabaci* biotypes on all host plants tested  
205 ranges from 30 to 33°C (Drost *et al.*, 1998). The developmental threshold for  
206 immatures belonging to the population studied here, estimated at 10.2°C, is lower  
207 than that reported in the literature, which ranges from 10.8°C (Von Arx *et al.*, 1983)  
208 to 12.5°C (Wang & Tsai, 1996).

209 Based on the range of temperatures tested here, survival rates of immature stages of  
210 *B. tabaci* from egg to adult are higher on the tomato cultivar Hilario than those  
211 reported on other cultivars of tomato (Tsai & Wang, 1996), Poinsettia (Enkegaard,  
212 1993) and cotton (Wagner, 1995). Different responses to extreme temperatures, i.e.  
213 in mortality of immatures, suggest that the Q-biotype population on tomato is more  
214 tolerant to high temperatures (>33°C) than diverse *B. tabaci* populations on  
215 aubergine (Wang & Tsai, 1996), cotton, and Poinsettia (Drost *et al.*, 1998).

216 Fecundity of *B. tabaci* is generally highly variable and depends on temperature  
217 (Enkegaard, 1993), host-plant species (Liu & Oetting, 1994), and cultivar (Navon *et al.*, 1991). Thus, the total number of eggs laid at 25°C by a Q-biotype female reared  
218 on tomato *cv Hilario* (94.2 eggs) is considerably lower than that reported in the  
219

220 literature for females reared on aubergine (223.67) (Wang & Tsai, 1996) and tomato  
 221 cv Suny Hybrid (165.55) (Tsai & Wang, 1996).  
 222 Wang & Tsai (1996) underline the importance of life table parameters to compare *B.*  
 223 *tabaci* populations and biotypes. On aubergine these authors (Wang & Tsai, 1996)  
 224 find a high intrinsic rate of increase at 25°C (0.192), 27°C (0.191), and 30°C (0.169)  
 225 compared with that found with the Q-biotype population on tomato at 25°C (0.106)  
 226 and 30°C (0.123). However, at 35°C, the net rate of increase of the B-biotype (0.073)  
 227 is lower than that of the French Q-biotype population (0.104). The tolerance to  
 228 extreme thermal conditions of the population newly introduced in France is also  
 229 confirmed by its shortest mean generation time recorded at 35°C (24.6 days).  
 230 The comparison between the results of this work and those from other studies (Wang  
 231 & Tsai, 1996; Muñoz & Nombela, 1996, 2001; Drost *et al.*, 1998; Nombela *et al.*,  
 232 2000, Muñoz, 2000) demonstrates clearly that the relationship between life-history  
 233 parameters and temperature is influenced highly by both insect biotype and host  
 234 plant species and variety. In spite of this high degree of variability within the *Bemisia*  
 235 complex, it is essential to better understand the population dynamics of the newly  
 236 introduced pest population in relation to temperature to improve control strategies  
 237 and evaluate its geographical extension capacity.

238

## 239 **References**

240

241 Arx, R. von, Baumgärtner, J. & Delucchi, V. (1983) A model to simulate the  
 242 population dynamics of *Bemisia tabaci* Genn (Homoptera: Aleyrodidae) on  
 243 cotton in the Sudan Gezira. *Zeitschrift für Angewandte Entomologie*, **96**, 341-  
 244 363.

- 245 Bethke J.A., Paine, T.D. & Nuessly, G.S. (1991) Comparative biology,  
246 morphometrics, and development of two populations of *Bemisia tabaci*  
247 (Homoptera: Aleyrodidae) on cotton and poinsettia. *Annals of Entomological*  
248 *Society of America*, **84**, 407-411.
- 249 Chermitti, B., Braham, M., Alonso, C., Beitia, F. & Cenis, J.L. (1997) Sur la  
250 présence en Tunisie des biotypes 'B' et 'non-B' de *Bemisia tabaci* (Homoptera:  
251 Aleyrodidae) et de leurs parasitoides associés. *IOBC/ WPRS Bulletin*, **20**, 108-  
252 113.
- 253 Claridge, M.F., Dawah, H.A. & Wilson, M.R. (1997) Species in insect herbivores  
254 and parasitoids – sibling species, host races and biotypes. *Species, the Units of*  
255 *Biodiversity*. (ed. by M.F., Claridge, H.A., Dawah, M.R., Wilson), pp. 247-272.  
256 Chapman & Hall, London.
- 257 Coudriet, D.L., Prabhaker, N., Kishaba, N. & Meyerdirk, D.E. (1985) Variation in  
258 development rate on different hosts and overwintering of the sweetpotato  
259 whitefly *Bemisia tabaci* (Homoptera: Aleyrodidae). *Environmental*  
260 *Entomology*, **14**, 516-519.
- 261 Curry, G.L. & Feldman, R.M. (1987) Mathematical foundations of population  
262 dynamics. Texas A& M University Press, College Station, 246 pp.
- 263 Dalmon, A., Cailly, M., Bouyer, S., Arnold-Gaulhiac, M., Cailly, A. & Goarant, G.  
264 (2003) Emergence de virus transmis par aleurodes dans les cultures de tomate en  
265 France. Proceedings of the International Symposium on Greenhouse Tomato:  
266 Integrated Crop Protection and Organic Production, CTIFL, Paris, 24-29.
- 267 De Barro, P.J., Driver, F., Trueman, J.W.H. & Curran, J. (2000) Phylogenic  
268 relationship of world populations of *Bemisia tabaci* (Gennadius) using ribosomal  
269 ITS1. *Molecular Phylogenetics and Evolution*, **16**, 29-36.

- 270 Demichelis, S., Bosco, D., Manino, A., Marian, D. & Caciagli, P. (2000) Distribution  
271 of *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotypes in Italy. *Canadian*  
272 *Entomologist*, **132**, 519-527.
- 273 Drost, Y.C., van Lenteren, J.C. & van Roermund, H.J.W. (1998) Life-history  
274 parameters of different biotypes of *Bemisia tabaci* (Hemiptera: Aleyrodidae) in  
275 relation to temperature and host plant: a selective review. *Bulletin of*  
276 *Entomological Research*, **88**, 219-229.
- 277 Enkegaard, A. (1993) The poinsettia strain of the cotton whitefly, *Bemisia tabaci*  
278 (Homoptera: Aleyrodidae), biological and demographic parameters on  
279 poinsettia (*Euphorbia pulcherima*) in relation to temperature. *Bulletin of*  
280 *Entomological Research*, **83**, 535-546.
- 281 Guirao, P., Beitia, F. & Cenis, J.C. (1997) Biotype determination of Spanish  
282 populations of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Bulletin of*  
283 *Entomological Research*, **87**, 587-593.
- 284 Horowitz, A.R., Denholm, I., Gorman, K., Cenis, J.L., Kontsedalov, S. & Ishaaya, I.  
285 (2003) Biotype Q of *Bemisia tabaci* identified in Israel. *Phytoparasitica*, **31**, 94-  
286 98.
- 287 Hulting, F.L., Orr, D.B. & Obrycki, J.J. (1990) A computer program for calculation  
288 and statistical comparison of intrinsic rates of increase and associated life table  
289 parameters. *Florida Entomologist*, **73**, 601-612.
- 290 Liu, T.X. & Oetting, R.D. (1994) Oviposition preference of *Bemisia tabaci*  
291 (Gennadius) on eight species of greenhouse-grown plants. *Journal of*  
292 *Agricultural Entomology*, **11**, 177-179.

293 Logan, J.A., Wollkind, D.J., Hoyt, S.C. & Tanigoshi, L.K. (1976) An analytic model  
 294 for description of temperature dependent rate phenomena in arthropods.  
 295 *Environmental Entomology*, **5**, 1133-1140.

296 Lopez-Avila, (1986) Natural enemies. *Bemisia tabaci – a literature survey on the*  
 297 *cotton whitefly with an annotated bibliography*, (ed. by M. J. Cock, M.), 121pp.  
 298 C.A.B. International Institute of Biological Control, Silwood Park, Ascot,  
 299 Berks.

300 Meyer, J.S., Ingersoll, C.G., McDonald, L.L. & Boyce, M S. (1986) Estimating  
 301 uncertainty in population growth rates: jackknife vs. bootstrap techniques.  
 302 *Ecology*, **67**, 1156-1166.

303 Monci, F., Navas-Castillo, J., Cenis, J.L., Lacasa, A., Benazoun, A. & Moriones, E.  
 304 (2000) Spread of tomato yellow leaf curl virus-Sar from the Mediterranean basin:  
 305 presence in the Canary Islands and Morocco. *Plant Disease*, **84**, 490

306 Muñoz, M. (2000) Host suitability of two biotypes of *Bemisia tabaci* on some  
 307 common weeds. *Entomologia Experimentalis et Applicata*, **95**, 63-70.

308 Muñoz, M. & Nombela, G. (1997) Development, oviposition and female longevity of  
 309 two biotypes of *Bemisia tabaci* (Homoptera: Aleyrodidae) on three varieties of  
 310 *Capsicum annuum* L. *IOBC/ WPRS Bulletin*, **20**, 143-146.

311 Muñoz, M. & Nombela, G. (2001) Differential variation in development of the B- and  
 312 Q-biotypes of *Bemisia tabaci* on sweet pepper *Capsicum annuum* L. at  
 313 constant temperatures. *Environmental Entomology*, **30**, 720-727.

314 Naranjo, S.E. (2001) Conservation and evaluation of natural enemies in IPM systems  
 315 for *Bemisia tabaci*. *Crop Protection*, **20**, 835-852.

- 316 Nava-Camberos, U., Riley, D.G. & Harris, M.K. (2001) Temperature and host plant  
317 effects on development, survival, and fecundity of *Bemisia argentifolii*  
318 (Homoptera: Aleyrodidae). *Environmental Entomology*, **30**, 55-63.
- 319 Navas-Castillo, J., Camero, R., Bueno, M. & Moriones, E. (2000) Severe yellowing  
320 outbreaks in tomato in Spain associated with infections of Tomato chlorosis  
321 virus. *Plant Disease*, **84**, 835-837.
- 322 Navon, A., Melamed, M.V., Zur, M. & Ben, M.E. (1991) Effects of cotton cultivars  
323 on feeding of *Heliothis armigera* and *Spodoptera littoralis* larvae and on  
324 oviposition of *Bemisia tabaci*. *Agriculture, Ecosystems and Environment*, **35**,  
325 73-80.
- 326 Oliviera, M.R.V., Henneberry, T.J. & Anderson, P. (2001) History, current status,  
327 and collaborative research projects for *Bemisia tabaci*. *Crop Protection*, **20**, 709-  
328 723.
- 329 Simón, B., Cenis, J.L., Demichelis, S., Rapisarda, C., Caciagli, P. & Bosco, D.  
330 (2003) Survey of *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotypes in Italy with  
331 the description of a new biotype (T) from *Euphorbia characias*. *Bulletin of*  
332 *Entomological Research*, **93**, 259-264.
- 333 Tsai, J.H. & Wang, K. (1996) Development and reproduction of *Bemisia argentifolii*  
334 on five host plants. *Environmental Entomology*, **25**, 810-816.
- 335 Van Lenteren, J.C. & Noldus, L.P.J.J. (1990) Whitefly-plant relationships:  
336 behavioral and ecological aspects. *Whiteflies: their bionomics, pest status and*  
337 *management*, (ed. by D Gerling), pp. 47-89. Intercept, Andover, Hampshire,  
338 U.K.

- 339 Wagner, T.L. (1995) Temperature-dependent development, mortality, and adult size  
340 of sweetpotato whitefly biotype B (*Homoptera: Aleyrodidae*) on cotton.  
341 *Environmental Entomology*, **24**, 1179-1188.
- 342 Wang, K. & Tsai, J.H. (1996) Development and reproduction of *Bemisia argentifolii*  
343 (Homoptera: Aleyrodidae) on five host plants. *Annals of Entomological Society*  
344 *of America*, **89**, 375-384.
- 345 Zalom, F.G., Castañe, C. & Gabarra, R. (1995) Selection of some winter-spring  
346 vegetable crop hosts by *Bemisia tabaci* (Homoptera: Aleyrodidae). *Journal of*  
347 *Economic Entomology*, **88**, 70-76.
- 348



**Table 1.** Developmental period (day  $\pm$  SE) of immature stages of *Bemisia tabaci* (Q biotype) at 5 constant temperatures.

Temp. (°C)	Stages						From egg to adult
	<i>n</i>	eggs	instar 1	instar 2	instar 3	instar 4	
17	80	21.5 $\pm$ 0.09a	9.4 $\pm$ 0.35a	7.5 $\pm$ 0.93a	8.0 $\pm$ 0.42a	9.4 $\pm$ 0.33a	55.8 $\pm$ 0.47a
21	80	14.0 $\pm$ 0.09b	7.1 $\pm$ 0.21b	4.1 $\pm$ 0.18b	8.8 $\pm$ 0.23a	5.6 $\pm$ 0.18b	39.6 $\pm$ 0.69b
25	80	10.4 $\pm$ 0.13c	4.3 $\pm$ 0.19c	3.7 $\pm$ 0.16bc	3.8 $\pm$ 0.15b	3.4 $\pm$ 0.12c	25.6 $\pm$ 0.26c
30	80	7.7 $\pm$ 0.06d	3.2 $\pm$ 0.13d	3.3 $\pm$ 0.15c	3.5 $\pm$ 0.15b	2.5 $\pm$ 0.09d	20.2 $\pm$ 0.24d
35	38	6.5 $\pm$ 0.10e	3.9 $\pm$ 0.17cd	3.3 $\pm$ 0.21c	3.5 $\pm$ 0.29b	3.3 $\pm$ 0.22c	20.5 $\pm$ 0.33d

Within columns means followed by the same letters are not significantly different ( $P>0.05$ ).

**Table 2.** Survivorship (percentage) of immature stages of *Bemisia tabaci* (Q biotype) at 5 constant temperatures.

Temp. (°C)	Stages							From egg to adult
	<i>n</i> eggs	egg	instar 1	instar 2	instar 3	instar 4	<i>n</i> adults	
17	80	91.3	84.9	96.8	73.3	63.6	38	48c
21	80	98.7	98.7	93.4	93.0	97.0	64	83a
25	80	100	93.8	94.7	98.6	97.1	68	85a
30	80	100	95.0	96.1	93.2	97.1	66	82a
35	38	100	97.4	89.2	78.8	92.3	24	63b

Within columns means followed by the same letters are not significantly different ( $P>0.05$ ).

**Table 3.** Fecundity (number of eggs per female  $\pm$  SE) and longevity (days  $\pm$  SE) of *Bemisia tabaci* (Q biotype) at 5 constant temperatures.

Temp. (°C)	<i>n</i>	Total fecundity	Longevity
17	30	49.3 $\pm$ 6.7b	39.6 $\pm$ 3.6a
21	30	105.3 $\pm$ 10.4a	27.3 $\pm$ 0.8b
25	30	94.2 $\pm$ 12.3a	21.9 $\pm$ 1.7c
30	30	58.6 $\pm$ 10.4b	14.6 $\pm$ 1.1d
35	30	41.0 $\pm$ 5.6b	8.5 $\pm$ 0.7e

Within columns means followed by the same letters are not significantly different ( $P>0.05$ ).

**Table 4.** Comparison of life table parameters of *Bemisia tabaci* (Q biotype) at 5 constant temperatures. n = number of females.  $r_m$  = Jackknife estimate of the intrinsic rate of increase. CI = confidence interval estimate of  $r_m$ . Ro = net reproductive rate (Standard error). G = mean generation time in day.  $\lambda$  = finite rate of increase =  $\exp(r_m)$ .

Temp. ( °C)	Parameters					
	<i>n</i>	$r_m$	95%CI	Ro	G	$\lambda$
17	30	0.045c	0.044-0.046	29.8 (0.2)	77.2	1.05
21	24	0.079b	0.078-0.080	52.5 (0.1)	49.9	1.08
25	18	0.106a	0.090-0.121	39.5 (5.5)	35.1	1.11
30	24	0.123a	0.103-0.142	23.6 (4.3)	26.5	1.13
35	23	0.104a	0.087-0.119	12.3 (1.8)	24.6	1.11

Within columns means followed by the same letters are not significantly different ( $P > 0.05$ ).

## Figure captions

**Fig. 1.** Influence of temperature on development time of *Bemisia tabaci* (Q biotype). Points: experimental values. curve simulated by Logan *et al.* (1976).


**Fig. 2.** Influence of temperature on survivorship of *Bemisia tabaci* (Q biotype). Points: experimental values. curve simulated by Curry & Feldman model (1987)

**Fig. 3.** Influence of temperature on female longevity of *Bemisia tabaci* (Q biotype). Points: experimental values. curve simulated by exponential model.


**Fig. 4.** Influence of temperature on fecundity of *Bemisia tabaci* (Q biotype). Points: experimental values. curve simulated by multiplicative exponential model.

**Fig. 5.** Influence of temperature on intrinsic rate of increase of *Bemisia tabaci* (Q biotype). Points: experimental values. curve: simulated by Logan *et al.* (1976) model.


**Fig. 1.**



**Fig. 2.**




**Fig. 3**





**Fig. 4.**



**Fig. 5.**

